Mon. Not. R. Astron. Soc. **000**, 000–000 (0000)    Printed 17 December 2012    (MN LaTeX style file v2.2)

arXiv:1212.2626v2 [astro-ph.SR] 14 Dec 2012

# A mean field dynamo from negative eddy diffusivity

Ebru Devlen$^{1,2\star}$, Axel Brandenburg$^{1,3}$ and Dhrubaditya Mitra$^{1}$
$^{1}$*Nordita, KTH Royal Institute of Technology and Stockholm University, Roslagstullsbacken 23, 10691 Stockholm, Sweden*
$^{2}$*Department of Astronomy & Space Sciences, Faculty of Science, University of Ege, Bornova 35100, Izmir, Turkey*
$^{3}$*Department of Astronomy, Stockholm University, 10691 Stockholm, Sweden*

17 December 2012, Revision: 1.34

## ABSTRACT

Using direct numerical simulations, we demonstrate that “flow IV” of Roberts (1972) exhibits dynamo action dominated by horizontally averaged large-scale magnetic field. With the test-field method we show that the magnetic eddy diffusivity is negative and overcomes the molecular diffusivity to explain this large-scale dynamo for magnetic Reynolds numbers above $\approx 8$. As expected for a dynamo of this type, the two horizontal field components grow independently of each other and have arbitrary amplitude ratios and phase differences. Small length scales of the mean magnetic field are shown to be stabilized by the eddy diffusivity becoming positive at wavenumbers larger than twice the characteristic wavenumber of the flow. For magnetic Reynolds numbers below $\approx 0.5$ the eddy diffusivity is confirmed to be positive, as expected for all incompressible flows. Earlier claims of a dynamo driven by a modified Taylor-Green flow through negative eddy diffusivity could not be confirmed.

**Key words:** Dynamo – magnetic fields – MHD – turbulence

## 1 INTRODUCTION

The equations of magnetohydrodynamics (MHD) permits growth of magnetic energy at the expense of kinetic energy. This phenomenon is called the dynamo effect (see e.g. Brandenburg & Subramanian 2005, for a recent review). If the dynamo effect gives rise to a magnetic field whose characteristic length scale is greater than that of the fluid, we call it a *large-scale* dynamo. Most astrophysical dynamos, including the solar dynamo and the galactic dynamo, are of this type.

To theoretically describe the large-scale dynamo one must average the equations of MHD over the small scales to write an effective equation for the large-scale magnetic field. This effective equation can be written down by using either mean-field theory (Steenbeck, Krause & Rädler 1966) or multiple-scale expansions (see e.g. Zheligovsky 2011, for a recent review). These equations contain turbulent transport coefficients: the $\alpha$ effect and turbulent (or eddy) diffusivity. In general, both are tensors whose complexity depends on the symmetries of the problem. Within the formalism of mean-field theory, it is generally a non-trivial task to calculate the turbulent transport coefficients even if we ignore the back-reaction of the magnetic field on the flow, i.e., for kinematic dynamos. For several kinematic problems, the turbulent transport coefficients have been calculated using the test-field method (Sur et al. 2008; Brandenburg et al. 2008a,b). Typically, it is found that the $\alpha$ effect gives rise to the growth of a large-scale magnetic field while the eddy diffusivity contributes to decay by effectively enhancing the molecular magnetic diffusivity. However, multiscale methods have shown that for certain flows the $\alpha$ effect can be zero, but the eddy diffusivity may turn out to be negative (see e.g., Lanotte et al. 1999; Zheligovsky et al. 2001). If, for such flows, the magnitude of the eddy diffusivity is greater than the molecular one, a large-scale dynamo can operate.

In this paper we first demonstrate, by direct numerical simulations (DNS) that for a particular flow (Roberts 1972), which will in the following be referred to as Roberts-IV flow, it is possible to drive a kinematic large-scale dynamo although the $\alpha$ effect (and the planar-averaged kinetic helicity) is indeed zero. Next, by using the test-field method, we show that such a dynamo can be accurately described by zero $\alpha$ effect but negative eddy diffusivity which dominates over molecular one.

## 2 THE ROBERTS-IV FLOW

In a remarkable paper, Roberts (1972) studied several two-dimensional periodic flow patterns that all produce dynamo action, which means that an infinitesimally weak magnetic field gets amplified exponentially in time. In connection with understanding the geodynamo, Tilgner (2004) studied in some detail the “flow IV” of Roberts (1972). We follow here Tilgner’s definition of the flow:

$$\boldsymbol{U} = u_0 \begin{pmatrix} \sqrt{2/f}\,\sin k_0x\cos k_0y \\ -\sqrt{2/f}\,\cos k_0x\sin k_0y \\ \sqrt{f}\,\sin k_0x \end{pmatrix}, \tag{1}$$

where $u_0$ characterizes the amplitude of the flow. Its vorticity, $\boldsymbol{W} = \boldsymbol{\nabla}\times\boldsymbol{U}$, is given by

---

$^\star$ E-mail:devlen@nordita.org

© 0000 RAS



**Figure 1.** Three components of the magnetic field on the periphery of the computational domain for $R_\mathrm{m}=20$, $f=1$, and domain size $L_x=L_y=L_z=2\pi/k_0$.

$$\boldsymbol{W}=u_0\begin{pmatrix}0\\ -\sqrt{f}\,k_0\cos k_0x\\ 2\sqrt{2/f}\,k_0\sin k_0x\sin k_0y\end{pmatrix}. \tag{2}$$

Here, the parameter $f$ determines the relative importance of eddy diffusion along the $z$ direction (large $f$) compared to that in the $xy$ plane (small $f$). The kinetic helicity density, $\boldsymbol{W}\cdot\boldsymbol{U}$, is given by

$$\boldsymbol{W}\cdot\boldsymbol{U}=\sqrt{2}\,u_0^2k_0\,(1+\sin^2k_0x)\sin k_0y \tag{3}$$

and is independent of $f$. Tilgner (2004) showed that in spite of the horizontally averaged kinetic helicity density, $\overline{\boldsymbol{W}\cdot\boldsymbol{U}}$, being zero; the Roberts IV flow gives rise to dynamo action. In other words, it leads to growing solutions of the induction equation,

$$\frac{\partial\boldsymbol{B}}{\partial t}=\boldsymbol{\nabla}\times(\boldsymbol{U}\times\boldsymbol{B}-\eta\boldsymbol{J}), \tag{4}$$

where $\eta$ is the microphysical (molecular) magnetic diffusivity, $\boldsymbol{B}$ is the magnetic field, $\boldsymbol{J}=\boldsymbol{\nabla}\times\boldsymbol{B}$ is the current density, and we have chosen our units such that the vacuum permeability is unity. Here and in the following, overbars denote $xy$ averaging.

Note, however, that Tilgner (2004) described the dynamo to be a small-scale one, i.e., the characteristic length scales of the magnetic field is of the same order as $1/k_0$. In the following, we obtain solutions to Equation (4) via DNS using the PENCIL CODE[1]. We do not evolve the flow, hence we study kinematic dynamo solutions. As an example of the resulting magnetic field, we show in Figure 1 the three components of the magnetic field at the periphery of the computational domain. It is remarkable that the resulting magnetic field has a large-scale component that survives horizontal averaging, i.e., $\overline{\boldsymbol{B}}=\overline{\boldsymbol{B}}(z,t)$ is non-vanishing; see Figure 2, where we show examples of the resulting mean field obtained by averaging the solution of the DNS. In other words, we have here an example of a mean-field dynamo, where $\overline{\boldsymbol{B}}$ is a solution of the horizontally averaged induction equation,

$$\frac{\partial\overline{\boldsymbol{B}}}{\partial t}=\boldsymbol{\nabla}\times\left(\overline{\boldsymbol{U}}\times\overline{\boldsymbol{B}}+\overline{\boldsymbol{\mathcal{E}}}-\eta\overline{\boldsymbol{J}}\right), \tag{5}$$

where $\overline{\boldsymbol{\mathcal{E}}}=\overline{\boldsymbol{u}\times\boldsymbol{b}}$ is the mean electromotive force resulting from correlations of fluctuating velocity and magnetic fields, $\boldsymbol{u}=\boldsymbol{U}-\overline{\boldsymbol{U}}$ and $\boldsymbol{b}=\boldsymbol{B}-\overline{\boldsymbol{B}}$, respectively. Empirically, we find that the horizontally averaged solutions of Equation (4) are of the form

$$\overline{\boldsymbol{B}}(z,t)=\begin{pmatrix}B_{0x}\cos(kz+\phi_x)\\ B_{0y}\cos(kz+\phi_y)\\ 0\end{pmatrix}e^{\lambda t}, \tag{6}$$

where $B_{0x}$, $B_{0y}$, $\phi_x$, and $\phi_y$ are arbitrary constants, i.e., the $x$ and $y$ components of the magnetic field evolve independently of each other and they have arbitrary phase shifts, depending just on the properties of the initial conditions; see Figure 2 for an example.



**Figure 2.** Examples of runs with different initial conditions showing the $x$ and $y$ components of the mean field (normalized by the rms value of the actual field) versus $z$, obtained from the DNS for $f=1$, $L_x=L_y=L_z=2\pi/k_0$, and $\eta=0.05\,u_0/k_0$, corresponding to $R_\mathrm{m}=20$.

Solutions of Equation (5) can be obtained by mean-field simulations (MFS) which requires a closed expression for $\overline{\boldsymbol{\mathcal{E}}}$ in terms of $\overline{\boldsymbol{U}}$ and $\overline{\boldsymbol{B}}$. This will be discussed in the following.

As pointed out by Tilgner (2004), the Roberts-IV flow has no $\alpha$ effect. This is also confirmed by our simulations. As this is a laminar flow, driving a dynamo via fluctuations of $\alpha$ effect (Mitra & Brandenburg 2012) is also not possible. This suggests that the observed mean field might be produced by a negative eddy diffusivity effect (Lanotte et al. 1999; Zheligovsky et al. 2001). To investigate such a possibility, we now apply the test-field method (TFM) of Schrinner et al. (2005, 2007); see also Brandenburg (2005); Brandenburg et al. (2008a,b) for its implementation in periodic geometry; to calculate the turbulent transport coefficients of the Roberts IV flow.

As is long recognized (Rädler 1976), the connection between $\overline{\boldsymbol{\mathcal{E}}}$ and $\overline{\boldsymbol{B}}$ is a nonlocal one that is described by a convolution of the form

$$\overline{\mathcal{E}}_i=\hat{\alpha}_{ij}\circ\overline{B}_j-\hat{\eta}_{ij}\circ\overline{J}_j, \tag{7}$$

where "$\circ$" denotes a convolution in space and time, i.e.,

$$\hat{\eta}_{ij}\circ\overline{J}_j=\int\!\!\int\hat{\eta}_{ij}(z-z',t-t')\,\overline{J}_j(z',t')\,\mathrm{d}z'\,\mathrm{d}t', \tag{8}$$

and likewise for $\hat{\alpha}_{ij}\circ\overline{B}_j$, but this term is vanishing for the Roberts-IV flow (Tilgner 2004). The hats on $\hat{\alpha}_{ij}$ and $\hat{\eta}_{ij}$ indicate that the corresponding quantities are integral kernels.

We emphasize that in Equation (7) we have made use of the fact that the only non-vanishing derivatives of a horizontally averaged mean field are $\partial\overline{B}_x/\partial z$ and $\partial\overline{B}_y/\partial z$, which and be expressed in terms of components of $\overline{\boldsymbol{J}}$, so the corresponding eddy diffusion tensor is just of rank 2, not, as in the general case of rank 3 (Krause & Rädler 1980).

In the TFM, the kernel formulation above is most naturally considered in Fourier space with

$$\tilde{\mathcal{E}}(k,\omega)=\tilde{\alpha}_{ij}(k,\omega)\tilde{B}_j(k,\omega)-\tilde{\eta}_{ij}(k,\omega)\tilde{J}_j(k,\omega), \tag{9}$$

---

[1] `http://pencil-code.googlecode.com/`

where tildes denote appropriately normalized Fourier transformation of the corresponding mean-field quantities (Brandenburg et al. 2008b; Hubbard & Brandenburg 2009).

In the following we consider a three-dimensional domain of size $L_x \times L_y \times L_z$. For most cases we choose cubic domains, i.e., $L_x = L_y = L_z = 2\pi/k_0$. We are interested in the special case of harmonic solutions of Equation (5) with fixed vertical wavenumber $k = k_0$ of a magnetic field that is growing or decaying exponentially proportional to $e^{\lambda t}$. So we are interested in the case $\omega = \mathrm{i}\lambda$. As an approximation, we begin by considering the case $\omega = 0$, i.e., we ignore the so-called memory effect; see Hubbard & Brandenburg (2009) for illustrating the departure in the case of the standard Roberts flow with helicity (also known as the Roberts-I flow).

## 3 RESULTS

### 3.1 Sign change of eddy diffusivity

As we have mentioned already, all components of $\tilde{\alpha}_{ij}(k,\omega)$ vanish for the Roberts-IV flow. Moreover, $\tilde{\eta}_{ij}(k,\omega)$ is isotropic, i.e., we can write $\tilde{\eta}_{ij} = \tilde{\eta}_\mathrm{t}\delta_{ij}$. The resulting values of $\tilde{\eta}_\mathrm{t}(k_0, 0)$ are shown in Figure 3 as a function of magnetic Reynolds number,

$$R_\mathrm{m} = u_0/\eta k_0. \tag{10}$$

For comparison with earlier work, we note that this definition of $R_\mathrm{m}$ is close to a definition in terms of the rms velocity of the flow (for $f = 1$ we have $u_\mathrm{rms} \approx 1.225\,u_0$) and the wavenumber of the energy-carrying eddies $k_\mathrm{f}$, i.e., $u_\mathrm{rms}/\eta k_\mathrm{f}$. If we approximate $k_\mathrm{f} \approx w_\mathrm{rms}/u_\mathrm{rms}$, where $w_\mathrm{rms}$ is the rms value of the fluctuating part of the vorticity, then we have $k_\mathrm{f} \approx 1.29\,k_0$. Therefore, we have $u_\mathrm{rms}/\eta k_\mathrm{f} \approx 0.95\,R_\mathrm{m}$, which is close to $R_\mathrm{m}$.

As is common to many turbulent transport coefficients (Sur et al. 2008; Brandenburg et al. 2008b), $\eta_\mathrm{t}$ grows linearly with $R_\mathrm{m}$ for $R_\mathrm{m} \lesssim 0.5$; see Figure 3. More importantly, $\eta_\mathrm{t}$ is positive, which is to be expected based on a calculation for incompressible flows using the second order correlation approximation (SOCA), which is valid for $R_\mathrm{m} \ll 1$. To show this, one uses the fact that the Fourier transform of the velocity correlation tensor is positive semidefinite. We note in passing that this is not true for potential flows, for which the eddy diffusion tensor is negative semidefinite for $R_\mathrm{m} \ll 1$; see Rädler et al. (2011) for a recent demonstration using the TFM.

Returning now to the Roberts-IV flow, which is indeed incompressible, we show in Figure 3 that $\eta_\mathrm{t}$ changes sign from positive to negative at $R_\mathrm{m} \approx 4$. This is clearly a result that cannot be recovered by SOCA. Moreover, for $R_\mathrm{m} \approx 8$, the total magnetic diffusivity, $\eta + \eta_\mathrm{t}(k, 0)$ becomes negative, i.e., dynamo action by the negative magnetic diffusivity effect is possible. The critical value of $R_\mathrm{m}$ agrees with that found above through DNS. The growth rate of the dynamo is given in implicit form as a solution of the equation

$$\lambda(k) = -[\eta + \eta_\mathrm{t}(k, \mathrm{i}\lambda)]k^2 \tag{11}$$

for $k = k_0$. However, it is common to approximate $\eta_\mathrm{t}(k_0, \omega)$ by $\eta_\mathrm{t}(k_0, 0)$, and we refer to the corresponding solution as

$$\lambda(k) \approx \lambda_\mathrm{disp}(k) = -[\eta + \eta_\mathrm{t}(k, 0)]k^2, \tag{12}$$

which is shown in the third panel of Figure 3 and compared with the growth rate $\lambda_\mathrm{DNS}$ obtained by solving Equation (4) through DNS. The agreement between $\lambda_\mathrm{disp}$ and $\lambda_\mathrm{DNS}$ is moderate and reminiscent of what has been found earlier (Hubbard & Brandenburg 2009). By using a test field that grows exponentially at a rate that is equal to the expected growth rate, $\lambda = \lambda_\mathrm{DNS}$, we find for $\eta = 0.020\,u_0/k_0$ (corresponding to $R_\mathrm{m} \approx 50$), the value $\eta_\mathrm{t}(k_0, \mathrm{i}\lambda) = -0.081\,u_0/k_0$ with $\lambda = 0.061\,u_0k_0$, instead of the value $-0.070\,u_0/k_0$ for $\lambda = 0$. Thus, perfect agreement between DNS and TFM is obtained once the memory effect included.



**Figure 3.** Turbulent magnetic diffusivity versus $R_\mathrm{m}$ for $f = 1$ and $k = k_0$.

### 3.2 Dependence on $f$

Let us now discuss the dependence on the parameter $f$, which characterizes the relative importance of vertical to horizontal eddy diffusion. We consider here the case of $R_\mathrm{m} = 20$ and $k = k_0$. The results are shown in Figure 4. The negative eddy diffusivity dynamo is found to be operating in the range $0.6 \leq f \leq 1.23$, i.e., when the vertical eddy diffusivity is not much larger than the horizontal.

As indicated by the dotted line in Figure 4, both for small and for large values of $f$, there is an approximate power law dependence with $|\eta_\mathrm{t}| \sim f^{1.7}$. However, in the range $1.3 < f < 3$ the TFM diverges and is unable to deliver useful results. Diverging results of the TFM are common and related to unstable eigenvalues of the associated homogeneous system of equations solved in the TFM. Usually, this problem can be avoided by focussing on the inhomogeneous term, which dominates the result at intermediate time intervals (Rheinhardt & Brandenburg 2010).



**Figure 4.** Dependence of eddy diffusivity on $f$ for $R_\mathrm{m} = 20$ and $k = k_0$. Negative values of $\eta_\mathrm{t}$ are indicated with open symbols, and the horizontal dash-dotted line indicates the region above which there is dynamo action when $\eta + \eta_\mathrm{t}$ is negative. The dotted line has a slope of 1.7 and is shown for orientation.



**Figure 5.** Wavenumber dependence of eddy diffusivity for $R_\mathrm{m} = 20$ and $f = 1$.

### 3.3 Scale dependence

Owing to the $k^2$ factor in Equations (11) and (12), one might expect dynamos driven by negative eddy diffusivity to grow faster at smaller scales, unless $\eta_\mathrm{t}$ becomes positive at larger values of $k$. In Figure 5 we show the $k$ dependence of $\eta_\mathrm{t}(k, 0)$. It turns out that $\eta_\mathrm{t}(k, 0)$ is approximately constant for $k \leq k_* \approx 1.125\,k_0$, and positive with an approximate dependence

$$\eta_\mathrm{t}(k,0) \approx \frac{0.31\,u_0/k_0}{(k-k_\infty)^{0.7}} \quad (\text{for } k - k_\infty), \tag{13}$$

where $k_\infty \approx 2.25\,k_0$. In the range $k_* < k < k_\infty$ we have two data points in Figure 5 that clearly deviate from the description above. In addition, there are several other values of $k$ in this range where the TFM again diverges and is unable to deliver useful results.

To illuminate the problem of intermediate $k$ values further, we now use DNS to compute the growth rate as a function of the domain size $L_z$, keeping $L_x = L_z = 2\pi/k_0$. The result is shown in Figure 6. It turns out that $\lambda$ ($= \lambda_\mathrm{DNS}$) has a maximum at $L_z k_0 \approx 3$. Thus, large scale separation, as assumed in some analytic studies (Lanotte et al. 1999), is neither needed nor necessarily helpful for the operation of this negative eddy diffusivity dynamo. Furthermore, for $L_z k_0 \leq 2.55$, no dynamo is possible and the field decays in an oscillatory fashion. The oscillation frequency $\omega$ grows sharply as $L_z$ decreases further; see the dotted line in Figure 6.



**Figure 6.** Dependence of the growth rate $\lambda$ and the oscillation frequency $\omega$ on the domain size $L_z$ for $R_\mathrm{m} = 20$ and $f = 1$. In all cases with $\lambda > 0$, we have $\omega = 0$.

## 4 MODIFIED TAYLOR–GREEN FLOWS

While the possibility of a dynamo driven by negative eddy diffusivity has not been previously quantified for the Roberts-IV flow, it was discussed in some detail by Lanotte et al. (1999) for the Taylor–Green (TG) flow,

$$\boldsymbol{U}_\mathrm{TG} = u_0 \begin{pmatrix} \sin k_0x \cos k_0y \cos k_0z \\ -\cos k_0x \sin k_0y \cos k_0z \\ 0 \end{pmatrix}, \tag{14}$$

and the modified TG flow, $\boldsymbol{U}_\mathrm{TG} + A\boldsymbol{U}_\mathrm{A} + B\boldsymbol{U}_\mathrm{B}$, where $A$ and $B$ denote the amplitudes of additional contributions proportional to

$$\boldsymbol{U}_\mathrm{A} = u_0 \begin{pmatrix} \sin 2k_0x \cos 2k_0z \\ \sin 2k_0y \cos 2k_0z \\ -(\cos 2k_0x + \cos 2k_0y) \sin 2k_0z \end{pmatrix}$$

and

$$\boldsymbol{U}_\mathrm{B} = u_0 \begin{pmatrix} (\sin k_0x \cos 3k_0y + \frac{5}{13} \sin 3k_0x \cos k_0y) \cos k_0z \\ -(\cos 3k_0x \sin k_0y + \frac{5}{13} \cos k_0x \sin 3k_0y) \cos k_0z \\ \frac{2}{13} (\cos k_0x \cos 3k_0y - \cos 3k_0x \cos k_0y) \sin k_0z \end{pmatrix},$$

respectively.

We perform kinematic DNS with this flow and we indeed find dynamo action. We carry out calculations for $\eta = 0.02\,u_0/k_0$, which corresponds to the case where the dynamo is mildly supercritical. We consider the following three cases: (a) A cube of size $L_x = L_y = L_z = 2\pi/k_0$, using $128^3$ meshpoints; (b) a cuboid with $L_z = 4\,L_x$ and $L_y = L_x = 2\pi/k_0$, using $128^2 \times 512$ meshpoints; and (c) a cuboid with $L_x = L_y = 4\,L_z$ and $L_z = 2\pi/k_0$. In all these three cases the large-scale field obtained by averaging over the $xy$ plane decays as a function of time, i.e., *no* large-scale dynamo is obtained. However, there still remains, in principle, the possibility of a large-scale field developing that is zero under Reynolds averaging but is non-zero under other averaging procedure, e.g., Fourier filtering. But even this possibility is ruled out because we observe no growth of a large-scale field at $k < k_0$;



**Figure 7.** Power spectra of DNS for the TG flow with $A = 1$ and $B = 0$ in a domain of size $L_x = L_y = L_z = 8\pi/k_0$ using $256^3$ mesh points and $\eta = 0.02\,u_0/k_0$ at different times during the exponential growth phase of the dynamo. The earliest time is shown as a dash-dotted line.

see Figure 7. This is found by calculating the spectrum of the magnetic field from simulations with $256^3$ meshpoints and a domain size of $L_x = L_y = L_z = 8\pi/k_0$. In other words, a dynamo is observed but it is *not* a large-scale dynamo. This is corroborated by the TFM which produces positive eddy diffusivity in all the above-mentioned cases. In case (a) $\eta_t = +0.135\,u_0/k_0$ for $A = 1$ and $B = 0$, and $\eta_t = +0.146\,u_0/k_0$ for $A = 1$ and $B = 1$; in case (b) $\eta_t = +0.158\,u_0/k_0$. However, in case (c) the TFM becomes unstable. These result are in complete contrast with the Roberts-IV flow, for which the corresponding values are of the order of unity; see Figure 2.

## 5 CONCLUSIONS

In the present work we have revisited the Roberts-IV flow using the TFM to compute the full set of turbulent transport coefficients. We confirm an earlier result of Tilgner (2004) that a dynamo is possible and that all components of the $\alpha$ tensor are vanishing. In addition, we find that there is a finite horizontally averaged mean magnetic field, which should be explicable in terms of mean-field dynamo theory. The TFM reveals that the eddy diffusivity tensor is diagonal. Moreover, in the regime where the dynamo is excited, the eddy diffusivity is sufficiently strongly negative such that the total (molecular plus eddy) diffusivity is negative. This is an unusual situation in that the horizontal components of the mean field are completely decoupled and grow independently with arbitrary relative amplitudes and phase shifts.

Many laminar flows are only slow dynamos,and the Roberts-IV flow is no exception. These dynamos are therefore not expected to be astrophysically relevant. However, the method used to study such dynamos (TFM combined with DNS) is now playing an important role in the study of astrophysical dynamos for turbulent flows. The present work highlights the accuracy of this method in that it enables us to pinpoint the detailed nature of a dynamo exhibiting a finite averaged magnetic field.

In the present case of laminar flow patterns, nonlocality is crucially important. In other words, turbulent transport is described by a convolution with the mean fields in space and time rather than just a multiplication. The TFM is particularly well suited to deal with such cases. For generic turbulent flows, as shown in earlier works by Hubbard & Brandenburg (2009) and Rheinhardt & Brandenburg (2012), we expect these transport kernels to have a relatively simple form and that complicated kernels, such as found here and in the earlier work (Rädler & Brandenburg 2009) are atypical. Note however, that even though most astrophysical flows are turbulent and are expected to become statistically homogeneous and isotropic at small scales; in practice large scale anisotropy and inhomogeneity plays an important role. In many of those cases nonlocality cannot be neglected and many Fourier modes need to be taken into account, as demonstrated by Chatterjee et al. (2011) for flows driven by the magnetic buoyancy instability.

## ACKNOWLEDGEMENTS

We thank Alessandra Lanotte for inspiring us to look for negative eddy diffusivity dynamos. Financial support from the Scientific & Technological Research Council of Turkey (TÜBİTAK) and the European Research Council under the AstroDyn Research Project 227952 are gratefully acknowledged. The computations have been carried out at the National Supercomputer Centre in Umeå and at the Center for Parallel Computers at the Royal Institute of Technology in Sweden.

## REFERENCES

Brandenburg A. 2005, Astron. Nachr., 326, 787
Brandenburg A., Subramanian K. 2005, Phys. Rep., 417, 1
Brandenburg A., Rädler K.-H., Rheinhardt M., Käpylä P. J. 2008a, ApJ, 676, 740
Brandenburg A., Rädler K.-H., Schrinner M. 2008b, A&A, 482, 739
Chatterjee P., Mitra D., Rheinhardt M., Brandenburg A. 2011, A&A, 534, A46
Hubbard A., Brandenburg A. 2009, ApJ, 706, 712
Krause F., Rädler K.-H. 1980, Mean-field Magnetohydrodynamics and Dynamo Theory (Oxford: Pergamon Press)
Lanotte A., Noullez A., Vergassola M., Wirth A. 1999, Geophys. Astrophys. Fluid Dyn., 91, 131
Mitra D., Brandenburg A., 2012, MNRAS, 420, 2170
Rädler K.-H. 1976, in Basic Mechanisms of Solar Activity, Proceedings from IAU Symposium No. **71** held in Prague, Czechoslovakia, ed. V. Bumba and J. Kleczek (D. Reidel Publishing Company Dordrecht), 323
Rädler K.-H., Brandenburg A. 2009, MNRAS, 393, 113
Rädler K.-H., Brandenburg A., Del Sordo F., Rheinhardt M. 2011, Phys. Rev. E, 84, 4
Rheinhardt, M., & Brandenburg, A. 2010, A&A, 520, A28
Rheinhardt M., Brandenburg A. 2012, Astron. Nachr., 333, 71
Roberts G. O. 1972, Phil. Trans. R. Soc., A 271, 411
Rogachevskii I., Kleeorin N. 2003, Phys. Rev. E, 68, 036301
Rogachevskii I., Kleeorin N. 2004, Phys. Rev. E, 70, 046310
Schrinner M., Rädler K.-H., Schmitt D., Rheinhardt M., Christensen U. 2005, Astron. Nachr., 326, 245
Schrinner M., Rädler K.-H., Schmitt D., Rheinhardt M., Christensen U. R. 2007, Geophys. Astrophys. Fluid Dyn., 101, 81
Steenbeck M., Krause F., Rädler K.-H. 1966, Z. Naturforsch., 21a, 369; see also the translation in Roberts & Stix, The turbulent dynamo, Tech. Note 60, NCAR, Boulder, Colorado (1971)

Sur S., Brandenburg A., Subramanian K. 2008, MNRAS, 385, L15

Tilgner A. 2004, Geophys. Astrophys. Fluid Dyn., 98, 225

Zheligovsky V. A., Podvigina O. M., Frisch U. 2001, Geophys. Astrophys. Fluid Dyn., 95, 227

Zheligovsky V. A., 2012, Large-Scale Perturbations of Magnetohydrodynamic Regimes, Springer Lecture Notes in Physics **829**, Springer, Berlin, 2011.

`$Header: /var/cvs/brandenb/tex/ebru/tilgner/paper.tex,v 1.34 2012-12-12 11:20:18 brandenb Exp $`